

FOUR NEW PLANETS ORBITING SOMEWHAT METAL-ENRICHED STARS¹

C.G. TINNEY², R. PAUL BUTLER³, GEOFFREY W. MARCY^{4,5}, HUGH R.A. JONES⁶, ALAN J. PENNY⁷, CHRIS MCCARTHY³, BRAD D. CARTER⁸, JADE BOND^{9,2}

Submitted to The Astrophysical Journal.

ABSTRACT

We report the detection of four new extra-solar planets from the Anglo-Australian Planet Search orbiting the somewhat metal-enriched stars HD 73526, HD 76700, HD 30177 and HD 2039. The planetary companion of HD 76700 has a circular orbit with a period of 3.98 d. With $M \sin i = 0.197 \pm 0.017 M_{\text{JUP}}$, or 0.69 times the mass of Saturn, is one of the lowest minimum mass extra-solar planets yet detected. The remaining planets all have elliptical orbits with periods ranging from 190.5 d to 4.4 yr. All four planets have been found orbiting stars from a sub-sample of twenty metal-enriched and faint ($V < 9$) stars, which was added to the Anglo-Australian Planet Search's magnitude-limited $V < 7.5$ main sample in October 1998. These stars were selected to be metal-enriched on the basis of their Strömgren photometry, and their enrichment has been subsequently confirmed by detailed spectroscopic analysis.

Subject headings: planetary systems – stars: individual (HD2039, HD73526, HD30177, HD76700)

1. INTRODUCTION

The Anglo-Australian Planet Search (AAPS) is a long-term planet detection program which aims to perform extra-solar planet detection and measurement at the highest possible precision. Together with programmes using similar techniques on the Lick 3 m and Keck I 10 m telescopes (Fischer et al. 2001; Vogt et al. 2000), it provides all-sky planet search coverage for inactive F, G, K and M dwarfs down to a magnitude limit of $V = 7.5$. Initial results from this programme (Tinney et al. 2001; Butler et al. 2001; Tinney et al. 2002a; Jones et al. 2002a; Butler et al. 2002a; Jones et al. 2002b) demonstrate that it achieves long-term, systematic velocity precisions of 3 m s^{-1} or better, for suitably stable stars, down to our main sample magnitude limit of $V < 7.5$.

AAPS is being carried out on the 3.9 m Anglo-Australian Telescope (AAT), using the University College London Echelle Spectrograph (UCLES) and an I₂ absorption cell. UCLES is operated in its 31 lines mm^{-1} mode. Prior to 2001 September, it was used with a MIT/LL 2048 $\times$ 4096 15 μm pixel CCD, and since then has been used with an EEV 2048 $\times$ 4096 13.5 μm pixel CCD. Our target sample includes 178 FGK stars with $\delta < -20^\circ$ and $V < 7.5$. Where age/activity information is available from R'_{HK} indices – see for example Henry et al. (1996); Tinney et al. (2002b) – we require target stars to have $\log R'_{HK} > -4.5$ (corresponding to ages greater than 3 Gyr). The observing and data processing procedures follow that described in Butler et al. (1996) and Butler et al. (2001).

In addition to our primary sample of $V < 7.5$ dwarfs, a

small sub-sample of twenty fainter dwarfs (down to $V < 9$) was added in October 1999, following suggestions that metal-enriched stars seemed to be preferentially revealing planets (see eg. Laughlin (2000) and references therein). These stars all had *uvby* photometry suggesting metal-enrichment. All stars in this fainter sub-sample were observed with a maximum exposure time of 300 s regardless of observing conditions. As a result, the velocity precisions achieved for these targets are not as high as those demonstrated for the AAPS main sample (Jones et al. 2002a; Butler et al. 2001).

2. CHARACTERISTICS OF THE HOST STARS

HD 30177 (HIP 21850, SAO 233633) is a chromospherically inactive ($\log R'_{HK} = -5.08$; Tinney et al. (2002b)) G8V star (Houck & Cowley 1975). HD 30177 was observed 105 times by the HIPPARCOS satellite and found to be photometrically stable with a standard deviation of 0.017 magnitudes. Its HIPPARCOS parallax of $18.3 \pm 0.8 \text{ mas}$ implies absolute magnitudes of $M_V = 4.72 \pm 0.09$ (ESA 1997) and $M_{bol} = 4.36 \pm 0.10$ (Cox 2000). Strömgren *uvby* photometry from the General Catalogue of Photometric Data¹⁰ (Hauck & Mermilliod 1997), together with the *uvby* calibrations of (Schuster & Nissen 1989) indicate a metallicity of $[\text{Fe}/\text{H}] = +0.20 \pm 0.16$. A detailed metallicity analysis has been performed for all the G-dwarfs in the AAPS (Bond 2002), which indicates a metallicity of $[\text{Fe}/\text{H}] = +0.19 \pm 0.09$, in agreement with the *uvby*-based estimate. The *b* – *y* colour of HD 30177 indicates $T_{eff} = 5320 \pm 20 \text{ K}$ (Hauck & Mermilliod 1997; Olsen

¹ Based on observations obtained at the Anglo-Australian Telescope, Siding Spring, Australia.

² Anglo-Australian Observatory, PO Box 296, Epping. 1710. Australia. cgt@aaoepp.aao.gov.au

³ Carnegie Institution of Washington, Department of Terrestrial Magnetism, 5241 Broad Branch Rd NW, Washington, DC 20015-1305

⁴ Department of Astronomy, University of California, Berkeley, CA, 94720

⁵ Department of Physics and Astronomy, San Francisco State University, San Francisco, CA 94132.

⁶ Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead CH41 1LD, UK

⁷ Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, U.K.

⁸ Faculty of Sciences, University of Southern Queensland, Toowoomba, 4350. Australia.

⁹ School of Physics, University of Sydney, Sydney, 2000. Australia.

¹⁰ Data obtained directly from <http://obswww.unige.ch/gcpd/gcpd.html>

1984). From HD30177's $\log R'_{HK} = -5.08$ we estimate intrinsic stellar velocity jitter to be in the range $2\text{--}4\text{ ms}^{-1}$ (Saar et al. 1998).

Figure 1 shows evolutionary tracks at $[\text{Fe}/\text{H}] = +0.2$ and $+0.1$ for stars of near-Solar mass from the compilation of Girardi et al. (2000). The $[\text{Fe}/\text{H}] = +0.1$ tracks were obtained by interpolation between the $[\text{Fe}/\text{H}] = +0.2$ and $[\text{Fe}/\text{H}] = 0.0$ tracks in Girardi et al. (2000). Based on these models and the metallicity measurements for HD 30177, its mass is estimated to be $0.95 \pm 0.05 M_{\odot}$, and it would appear (in common with HD 73526 and HD 76700) to be beginning its evolution off the main sequence.

HD 73526 (HIP 42282, SAO 220191) is a G6V dwarf (Houck 1978) for which no R'_{HK} estimate is currently available. It was observed 137 times by HIPPARCOS, but its relative faintness means that only a 0.02 magnitude standard deviation limit is placed on its photometric stability. Its HIPPARCOS parallax is $10.6 \pm 1.0\text{ mas}$, which indicates $M_V = 4.1 \pm 0.2$ and $M_{bol} = 3.7 \pm 0.2$ (ESA 1997; Cox 2000). As for HD 30177 (see above), Strömgren *uvby* photometry indicates a metallicity of $[\text{Fe}/\text{H}] = +0.10 \pm 0.16$ (Hauck & Mermilliod 1997; Schuster & Nissen 1989), while a detailed metallicity analysis gives $[\text{Fe}/\text{H}] = +0.11 \pm 0.10$ (Bond 2002). The GCPD $b - y$ colour of HD 73526 indicates $T_{eff} = 5450 \pm 20\text{ K}$ (Hauck & Mermilliod 1997; Olsen 1984). Based on the evolutionary tracks shown in Figure 1, HD 73526 is estimated to have a mass of $1.05 \pm 0.05 M_{\odot}$.

HD 76700 (HIP 43686, SAO 250370, LTT 3291) is catalogued as a G6V dwarf by SIMBAD, as G5 by the Henry-Draper catalogue (Cannon & Pickering 1918-24) and as G8V by HIPPARCOS (ESA 1997). The HIPPARCOS B-V colour ($+0.745$) would indicate a G8 spectral type is more likely than G5 or G6, though the GCPD $b - y$ colour (Hauck & Mermilliod 1997) would indicate a G6 type is more appropriate. No R'_{HK} estimate is available. It was observed 109 times by HIPPARCOS, which found it to have constant magnitude with a standard deviation of 0.012 magnitude. Its HIPPARCOS parallax is $16.8 \pm 0.7\text{ mas}$, which indicates $M_V = 4.3 \pm 0.1$ and $M_{bol} = 3.9 \pm 0.2$ (ESA 1997; Lang 1992). Strömgren *uvby* photometry indicates a metallicity of $[\text{Fe}/\text{H}] = +0.14 \pm 0.16$ (Hauck & Mermilliod 1997; Schuster & Nissen 1989), while a detailed metallicity analysis gives $[\text{Fe}/\text{H}] = +0.10 \pm 0.11$ (Bond 2002). The GCPD $b - y$ colour indicates a $T_{eff} = 5423 \pm 20\text{ K}$ (Hauck & Mermilliod 1997; Olsen 1984). Based on the evolutionary tracks shown in Figure 1, HD 76700 is estimated to have a mass of $1.00 \pm 0.05 M_{\odot}$. There is no published $\log R'_{HK}$ measurement for HD 76700. Because the velocity amplitude observed in this star is small ($26 \pm 2\text{ ms}^{-1}$), a Ca HK spectrum was acquired to determine the level at which our results could be affected by velocity jitter. Figure 2 shows this spectrum for HD 76700 compared to those for several objects of similar spectral type, along with their spectral types and $\log R'_{HK}$ values (Tinney et al. 2002b). HD 76700 shows no evidence for a line reversal. Based on this comparison we assign an upper limit to the $\log R'_{HK}$ for HD 76700 of -4.9 , from which we estimate its intrinsic stellar velocity jitter to be in the range $3\text{--}6\text{ ms}^{-1}$ or less (Saar et al. 1998).

HD 2039 (HIP 1931, SAO 23205) is a chromospherically

inactive ($R'_{HK} = -4.91$; Tinney et al. (2002b)) star classified as G2/G3 IV/V by Houck & Cowley (1975). It was observed 147 times by HIPPARCOS, but its relative faintness means that only a 0.022 magnitude standard deviation limit is placed on its photometric stability. Its HIPPARCOS parallax is $11.1 \pm 1.1\text{ mas}$, which indicates $M_V = 4.25 \pm 0.24$ and $M_{bol} = 4.22 \pm 0.25$ (ESA 1997; Cox 2000). There are multiple measures of this star's Strömgren *uvby* photometry, the mean of which indicate a metallicity of $[\text{Fe}/\text{H}] = +0.10 \pm 0.16$ (Hauck & Mermilliod 1997; Schuster & Nissen 1989). A detailed metallicity analysis gives $[\text{Fe}/\text{H}] = +0.10 \pm 0.11$ (Bond 2002). The $b - y$ colour of HD 2039 indicates $T_{eff} = 5675 \pm 20\text{ K}$ (Hauck & Mermilliod 1997; Olsen 1984). Based on the evolutionary tracks shown in Figure 1, HD 2039 is estimated to have a mass of $0.98 \pm 0.05 M_{\odot}$.

3. RADIAL VELOCITY OBSERVATIONS AND ORBITAL SOLUTIONS

Fifteen observations of HD 30177 are listed in Table 1. The column labelled "Uncertainty" is the velocity uncertainty produced by our least-squares fitting procedure. This fit simultaneously determines the Doppler shift and the spectrograph point-spread function (PSF) for each observation made through the iodine cell, given an iodine absorption spectrum and an "iodine free" template spectrum of the object (Butler et al. 1996). The uncertainty is derived for each measurement by taking the mean of four hundred useful spectral regions (each $2\text{ \AA}$ long) from each exposure. This uncertainty includes the effects of photon-counting uncertainties, residual errors in the spectrograph PSF model, and variation in the underlying spectrum between the template and "iodine" epochs. All velocities are measured relative to the zero-point defined by the template observation. Only observations where the uncertainty is less than twice the median uncertainty are listed. These data are shown in Figure 3. The figure shows the best-fit Keplerian model for the data, with the resultant orbital parameters listed in Table 2. Due to HD 30177's faintness compared to the AAPS main sample, the residuals about the fit (14 ms^{-1}) are significantly higher than the baseline 3 ms^{-1} precision level demonstrated for the main sample (Butler et al. 2001; Jones et al. 2002a).

Eighteen observations of HD 73526 are listed in Table 3, and they are shown in Figure 4 along with a Keplerian fit to the data with the orbital parameters listed in Table 2. Note again the larger residuals (17 ms^{-1}) compared to the AAPS main sample. Twenty-four observations of HD 76700 are listed in Table 4, and they are shown in Figure 5 along with a Keplerian fit to the data with the orbital parameters listed in Table 2. The thirty-six observations of HD 2039 are listed in Table 5, and they are shown in Figure 5 along with a Keplerian fit to the data with the orbital parameters listed in Table 2.

4. DISCUSSION

The resultant minimum companion mass for HD 76700 is $M \sin i = 0.197 \pm 0.017 M_{JUP}$, with an orbital semi-major axis $a = 0.049 \pm 0.004\text{ au}$ and eccentricity $e = 0.00 \pm 0.04$. This zero eccentricity is consistent with the expectation that a planet with a period of just $3.971 \pm 0.001\text{ d}$ will almost certainly lie in an orbit which has been tidally

circularised (Marcy & Butler 1998). The resulting orbital parameters for HD 76700 place it amongst the planetary companions with the lowest known minimum masses. HD 76700, joins HD 49674, HD 16141, HD 168746 & HD 46375 (Butler et al. 2002b; Marcy, Butler & Vogt 2000; Pepe et al. 2002) in the group of extra-solar planetary companions with measured minimum masses less than a Saturn mass ($0.299 M_{\text{JUP}}$).

The remaining three extra-solar planets all have elliptical orbits, though the ellipticity of the orbit for the companion to HD 30177 ($e=0.22\pm0.17$) is not different from zero with great significance. As further data are acquired over the coming years this parameter will become far better constrained, and it is possible that this extra-solar planet could turn out to be in a substantially circular orbit. If so this system would join with the other known nearly circular systems with gas-giant planets lying in orbits between where the Earth and Jupiter lie in our own Solar System (ϵ Ret, HD 4208, the outer components of 47 UMa, HD 28185 (Jones et al. 2002a)). These extra-solar planets would seem to indicate that gas-giants exist in nearly circular orbits with semi-major axes all the way out to, and beyond that of Jupiter, as confirmed recently by the detection of the outer planet in the 55 Cnc system (Marcy et al. 2002a).

Of the twenty metal-enriched stars included as a sub-sample along with our main sample in late 1999, five have revealed the presence of planetary companions (the four planets discussed here – HD 30177, 73526, 76700 and 2039 – along with the previously known companion to HD 83443 (Butler et al. 2002a; Mayor et al. 2002)). This gives us a lower limit (there may be longer period or lower mass planets present which we cannot yet detect) to the discovery rate of $25\pm11\%$ for this “metallicity-biased” sub-sample. This compares with the overall discovery rates estimated for the Keck, Lick and AAPS of $\sim 8\%$ (ie 8% of stars sur-

veyed have planets in orbits within 3.5 a.u. of their host stars (Butler et al. 2002c)) – a difference which, while not of great statistical significance, is not unexpected given that extra-solar planets seem to be being found preferentially around metal-enriched stars (eg. Reid (2002); Laughlin (2000) and references therein). It is also interesting to note (even if perhaps not statistically significant) that all four of the stars in this paper would seem to be beginning their evolution off the main sequence (see Fig 1).

These results would suggest that the biasing of planet surveys toward metal-enriched host stars *may* offer a benefit in the planet detection rate. However, such an increased discovery rate must be balanced against the fact that it will produce an inherently biased sample of extra-solar planets. With the total number of extra-solar planets still numbering less than a hundred, and the parameters of this ensemble of planets still poorly placed in a scheme of extra-solar planetary formation and evolution, now is not the time for planet searches to begin biasing their large surveys in the chase for better “hit rates” at the expense of scientific utility.

The Anglo-Australian Planet Search team would like to gratefully acknowledge the support of Dr Brian Boyle, Director of the AAO, the contributions and assistance of Drs Stuart Ryder and Debra Fischer, and the superb technical support which has been received throughout the programme from AAT staff - in particular R.Patterson, D.Stafford, F.Freeman, S.Lee, J.Pogson, G.Kitley and J. Stevenson. We further acknowledge support by; the partners of the Anglo-Australian Telescope Agreement (CGT,HRAJ,AJP); NASA grant NAG5-8299 & NSF grant AST95-20443 (GWM); NSF grant AST-9988087 (RPB); and Sun Microsystems. NSO/Kitt Peak FTS data used here were produced by NSF/NOAO.

REFERENCES

- Alonso, A., Arribas, S. & Martínez-Roger, C. 1995, *A&A*, 297, 197
 Cox, A.N. (ed.) 2000, *Allen's Astrophysical Quantities*, Springer: New York
 Bond, J. et al. (2002), in preparation.
 Butler, R.P., Marcy, G.W., Williams, E., McCarthy, C. & Dosanjh, P. 1996, *PASP*, 108, 500
 Butler, R.P., Tinney, C.G., Marcy, G.W., Jones, H.R.A., Penny, A.J., Apps, K. 2001, *ApJ*, 555, 410
 Butler, R.P. et al. 2002a, *ApJ*, in press (astro-ph/0206178)
 Butler, R.P., Marcy, G., Vogt, S., Fischer, D., Henry, G., Laughlin, G. & Wright 2002b, *ApJ*, submitted (<http://exoplanets.org/esp/hd49674/hd49674.shtml>)
 Butler, R.P., Vogt, S.S., Tinney, C.G., Jones, H.R.A., Penny, A.J., Apps, K. 2002c, in *Planetary Systems in the Universe: Observation, Formation and Evolution*, IAU Symp. 202, ed. A.J.Penny, P.Artymowicz, A.M. Lagrange & S.Russel, in press (<http://exoplanets.org/stats2000.html>)
 Cannon, A.J. & Pickering, E.C (1918-1924), *The Henry Draper Catalogue and Extension*, Harv. Ann. 91-100
 ESA, 1997, *The HIPPARCOS and Tycho Catalogues*, ESA SP-1200
 Fischer, D. A., Marcy, G. W., Butler, R. P., Vogt, S. S., Frink, S., & Apps, K. 2001, *ApJ*, 551, 1107
 Giradri, L., Bressan, A., Bertelli, G. & Chiosi, C. 2000 *A&AS*, 141, 371
 Hauck, B. & Mermilliod, M. 1997, *A&AS*, 124, 349
 Hauck, B. & Mermilliod, M. 1998, *A&AS*, 129, 431
 Henry, T.J., Soderblom, D.R., Donahue, R.A. & Baliunas, S.L. 1996, *AJ*, 111, 439.
 Houck, N. and Cowley, A.P. 1975, *Michigan Catalogue of Two Dimensional Spectral Types for the HD stars*, Volume 1, Michigan Spectral Survey: Ann Arbor
 Houck, N. 1978, *Michigan Catalogue of two dimensional spectral types for the HD stars*, Volume 2, Michigan Spectral Survey: Ann Arbor
 Jones, H.R.A., Butler, R.P., Marcy, G.W., Tinney, C.G., Penny, A.J., McCarthy, C. & Carter, B.D. 2002, *MNRAS*, in press (astro-ph/0112084)
 Jones, H.R.A., Butler, R.P., Marcy, G.W., Tinney, C.G., Penny, A.J., McCarthy, C., Carter, B.D. & Pourbaix, D. 2002, *MNRAS*, in press (astro-ph/0206216)
 Kurucz, R.L., Furenlid, I. Brault, J. & Testerman, L. 1984, *Solar Flux Atlas from 296 to 1300 nm*, National Solar Observatory Atlas No. 1
 Lang, K.R. 1992, *Astrophysical Data: Planets and Stars*, Springer-Verlag: New York
 Laughlin, G. 2000, *ApJ*, 545, 1064
 Marcy, G.W. & Butler, R.P. 1998, *ARA&A*, 36, 57
 Marcy, G.W., Butler, R.P. & Vogt, S.S. 2000, *ApJ*, 536, L43
 Marcy, G.W., Butler, R.P., Fischer, D.A., Laughlin, G., Vogt, S.S., Henry, G.W. & Pourbaix, D. 2002, *ApJ*, submitted (<http://exoplanets.org/esp/55Cnc/55Cnc.shtml>)
 Mayor, M., Naef, D., Pepe, F., Queloz, D., Santos, N.C., Udry, S. & Burnet, M. 2002, in *Planetary Systems in the Universe: Observation, Formation and Evolution*, IAU Symp. 202, ed. A.J.Penny, P.Artymowicz, A.M. Lagrange & S.Russel, in press (http://obswww.unige.ch/Preprints/Preprints/iau202_mayor.ps.gz)
 Olsen, E.H. 1984, *A&AS*, 57, 443
 Pepe, F. et al. 2002, *A&A*, 388, 632
 Reid, I.N. *PASP*, 114, 306
 Saar, S.H., Marcy, G.W. & Butler, R.P. 1998, *ApJ*, 498, L153
 Schuster, W.J. & Nissen, P.E. 1989, *A&A*, 221, 65

Tinney, C.G., Butler, R.P., Marcy, G.W., Jones, H.R.A., Penny, A.J., Vogt, S.S., Apps, K. & Henry, G.W. 2001, ApJ, 551, 507
 Tinney, C.G., Butler, R.P., Marcy, G.W., Jones, H.R.A., Penny, A.J., McCarthy, C. & Carter, B.D. 2001, ApJ, 571, 528

Tinney, C.G., McCarthy, C., Jones, H.R.A., Butler, R.P., Carter, B.D., Marcy, G.W., & Penny, A.J. 2002, MNRAS, 332, 759
 Vogt, S.S., Marcy, G.W., Butler, R.P. & Apps, K. 2000, ApJ, 536, 902




FIG. 1.— Evolutionary tracks from the compilation of Girardi et al. (2000). The Zero Age Main Sequence (ZAMS) is shown as a solid line, with evolutionary models at the indicated masses shown as *dot-dashed* lines. The open circles and uncertainties indicate the stars under discussion in this paper, with their estimated uncertainties indicated in brackets. The $[\text{Fe}/\text{H}] = +0.2$ tracks are directly from Girardi et al. (2000), while the $[\text{Fe}/\text{H}] = +0.1$ tracks were obtained by interpolation between the catalogued $[\text{Fe}/\text{H}] = +0.2$ and 0.0 models.




FIG. 2.— UCLES spectrum in the region of the Ca H line for HD 76700 and several comparison objects with $\log R'_{HK}$ measurements from Tinney et al. (2002b). For each object the spectral type and measured $\log R'_{HK}$ value is shown. Based on these we assign an upper limit to the $\log R'_{HK}$ value for HD 76700 of -4.9.




FIG. 3.— AAT Doppler velocities for HD 30177 from 1998 November to 2002 May. The solid line is a best fit Keplerian with the parameters shown in Table 2. The rms of the velocities about the fit is 14 m s^{-1} . Assuming $0.95 M_{\odot}$ for the primary, the minimum ($M \sin i$) mass of the companion is $7.7 \pm 1.5 M_{\text{JUP}}$, and the semi-major axis is $2.6 \pm 0.9 \text{ a.u.}$




FIG. 4.— AAT Doppler velocities for HD 73526 from 1999 January to 2002 June. The solid line is a best fit Keplerian with the parameters shown in Table 2. The rms of the velocities about the fit is 18 m s^{-1} . Assuming $1.05 M_{\odot}$ for the primary, the minimum ($M \sin i$) mass of the companion is $3.0 \pm 0.3 M_{\text{JUP}}$, and the semi-major axis is $0.66 \pm 0.05 \text{ a.u.}$




FIG. 5.— AAT Doppler velocities for HD 76700 from 1999 February to 2002 June, phased at the best fit Keplerian period of 3.971 d (*left panel*) and plotted unphased from the last three observing runs in 2002 (*right panel*). The plotted Keplerian has the parameters shown in Table 2. The rms of the velocities about the fit is 6.2 m s^{-1} . Assuming $1.0 M_{\odot}$ for the primary, the minimum ($M \sin i$) mass of the companion is $0.197 \pm 0.017 M_{\text{JUP}}$, and the semi-major axis is $0.049 \pm 0.004 \text{ a.u.}$




FIG. 6.— AAT Doppler velocities for HD 2039 from 1998 November to 2002 June, with the best fit Keplerian with parameters shown in Table 2. The rms of the velocities about the fit is 15 m s^{-1} . Assuming $0.98 M_{\odot}$ for the primary, the minimum ($M \sin i$) mass of the companion is $5.1 \pm 1.7 M_{\text{JUP}}$, and the semi-major axis is $2.2 \pm 0.2 \text{ a.u.}$

TABLE 1
VELOCITIES FOR HD 30177

JD ^a (-2451000)	RV ^a (m s ⁻¹)	Uncertainty (m s ⁻¹)
118.0974	77.9	10.9
119.1924	37.2	17.1
121.1514	49.7	14.8
157.1022	95.4	8.6
211.9834	92.4	11.3
212.9660	92.5	10.4
213.9995	92.3	9.1
214.9506	104.0	9.3
525.9973	26.9	6.9
630.9156	-0.5	9.9
768.3296	-61.5	10.2
921.1075	-117.8	10.2
1127.3205	-168.5	12.4
1188.2532	-172.9	7.8
1358.9181	-173.4	7.8

^aJulian Dates (JD) are barycentric. Radial Velocities (RV) are barycentric, but have an arbitrary zero-point determined by the radial velocity of the template, as described in Section 3

TABLE 2
ORBITAL PARAMETERS

Parameter	HD 30177	HD 73526	HD 76700	HD 2039
Orbital period P (d)	1620 $\pm$ 800	190.5 $\pm$ 3.0	3.971 $\pm$ 0.001	1190 $\pm$ 150
Velocity amp. K (m s $^{-1}$)	140 $\pm$ 10	108 $\pm$ 8	25 $\pm$ 2	136 $\pm$ 30
Eccentricity e	0.22 $\pm$ 0.17	0.34 $\pm$ 0.08	0.00 $\pm$ 0.04	0.69 $\pm$ 0.15
ω ($^{\circ}$)	288 $\pm$ 40	207 $\pm$ 30	-	333 $\pm$ 20
$a_1 \sin i$ (km)	(3.0 $\pm$ 1.5) $\times 10^6$	(0.265 $\pm$ 0.01) $\times 10^6$	1381 $\pm$ 1.2	(1.61 $\pm$ 0.45) $\times 10^6$
Periastron Time (JD-245000)	1027 $\pm$ 200	951 $\pm$ 12	1212.9 $\pm$ 0.1	836 $\pm$ 150
$M \sin i$ (M _{JUP})	7.7 $\pm$ 1.5	3.0 $\pm$ 0.3	0.197 $\pm$ 0.017	5.1 $\pm$ 1.7
a (AU)	2.6 $\pm$ 0.9	0.66 $\pm$ 0.05	0.049 $\pm$ 0.004	2.2 $\pm$ 0.2
RMS about fit (m s $^{-1}$)	14	18	6.2	15

TABLE 3
VELOCITIES FOR HD 73526

JD ^a (-2451000)	RV ^a (m s $^{-1}$)	Uncertainty (m s $^{-1}$)
212.1302	23.5	12.9
213.1314	66.8	13.7
214.2389	61.6	14.9
236.1465	48.4	10.9
630.0280	35.5	11.0
717.9000	-120.7	12.8
920.1419	-32.0	13.4
984.0378	78.0	9.3
1009.0976	71.8	9.7
1060.8844	-58.9	7.4
1091.8465	-149.8	13.6
1386.9003	51.8	6.8
1387.8921	64.0	6.2
1420.9248	-14.4	7.8
1421.9199	-9.6	6.5
1422.8602	-9.8	7.3
1424.9237	-19.1	10.4
1454.8529	-109.0	6.3

^aAs for Table 1

TABLE 4
VELOCITIES FOR HD 76700

JD ^a (-2451000)	RV ^a (m s ⁻¹)	Uncertainty (m s ⁻¹)
212.1565	9.2	6.7
213.1501	40.3	7.6
214.2583	-21.3	8.2
274.0177	-18.6	7.9
530.1791	26.7	8.8
683.8938	-19.8	5.7
920.1606	29.7	7.6
984.0068	22.8	7.1
1009.0638	-26.8	7.5
1060.9036	-24.3	4.7
1091.8517	-16.8	7.4
1129.8425	18.2	6.7
1359.0760	-14.3	6.6
1386.9145	-20.0	3.8
1387.9170	23.2	3.9
1388.9898	16.5	3.4
1389.8741	-11.8	3.9
1420.9418	7.3	4.5
1421.9331	-19.3	3.1
1422.8793	-11.0	2.7
1424.9439	0.6	5.3
1452.8943	-4.2	3.8
1454.8682	-3.2	3.9
1455.8565	22.8	3.1

^aAs for Table 1

TABLE 5
VELOCITIES FOR HD 2039

JD ^a (-2451000)	RV ^a (m s ⁻¹)	Uncertainty (m s ⁻¹)
118.0578	24.0	8.4
118.9610	-23.8	16.9
119.9445	0.0	11.5
121.0385	-33.0	16.1
211.9514	-10.1	18.7
212.9234	3.9	13.3
213.9749	-3.9	17.0
214.9171	-14.1	13.2
386.3227	-54.7	17.1
387.2981	-33.2	15.3
411.2293	-14.2	15.6
414.2585	-16.5	10.6
473.0883	-42.5	10.6
525.9286	-78.7	13.9
527.9226	-28.9	11.0
745.2702	-51.8	17.4
828.0703	-43.1	12.5
828.9944	-49.3	10.4
829.9757	-45.1	11.7
856.0702	-33.5	14.6
919.9434	-25.7	14.7
920.9672	-22.5	14.5
1093.2947	136.2	14.4
1127.2341	98.9	15.1
1151.2230	65.7	8.4
1152.0860	66.4	9.2
1154.2124	74.6	11.4
1187.0957	55.1	9.7
1188.0300	48.7	8.4
1189.1502	58.6	11.6
1190.0932	48.5	8.4
1422.3281	-9.7	11.2
1425.3322	-3.2	7.7
1455.2853	-29.4	6.1

^aAs for Table 1